

## Assessment of Bio-Treatments on Growth Performance of *Emblica officinalis* in Nursery

Anubha Srivastav<sup>1</sup>, Rambir Singh<sup>2</sup>, Anita Tomar<sup>3</sup>, Hari Om Shukla<sup>4</sup> & Yogeshkumar Agarwal<sup>5</sup>

<sup>1, 3, 4</sup> Forest Research Centre for Eco-rehabilitation, Prayagraj

<sup>2</sup> Forest Research Institute, Dehradun

<sup>5</sup> College of Forestry, SHUATS, Prayagraj

Corresponding Mail: [anubhasri\\_csfer@icfre.org](mailto:anubhasri_csfer@icfre.org)

### Abstract

The experiments were conducted on “Assessment of Bio-treatments on Growth Performance of *Emblica officinalis* in Nursery”. At Forest research center for Eco-Rehabilitation. *Emblica officinalis* were graded and uniform size was used for raising seedlings. The seedlings were raised in 13 × 26 cm. size polyethylene bags with a potting mixture of unsterilized sand: loam soil: farm yard manure (1:1:1). After germination in polyethylene bags 10 ml / g of broth / culture of 11 bio-treatments were inoculated in the polybags by making holes in the rhizosphere of seedlings. The treatments taken for the study were as following: T<sub>1</sub>-Control, T<sub>2</sub>-Phosphobacterium, T<sub>3</sub>-Bio P Plus, T<sub>4</sub>-Azotobacter, T<sub>5</sub>-Bio-K-Plus, T<sub>6</sub>-Bio-N-Plus, T<sub>7</sub>- JivaAmrit-liquid, T<sub>8</sub>- JivaAmrit-powder, T<sub>9</sub>-Vermicompost, T<sub>10</sub>-Azoline, T<sub>11</sub>-Rhizogold, and T<sub>12</sub>-FYM. The experiment was set up in a completely randomized block design with 12 treatments and three replications. Results revealed that the treatments T<sub>4</sub>- Azotobacter, T<sub>6</sub> - Bio-N-Plus, T<sub>7</sub>-Jiva Amrit-liquid, T<sub>8</sub>-Jiva Amrit-powder, and T<sub>9</sub>-Vermicompost remarkably showed very good increment in height and root length as compared to control with 11.53, 10.43, 10.97, 12.97 & 11.07 cm and 20.17, 21.27, 22.50, 20.63 & 22.27 cm respectively. Similarly, the survival was also recorded after 90 days of experimentation. It clear from the data that for most of the treatments, survival was 100 % but for control, it was only 80 %. Successful functioning of introduced microbial bio-inoculants influenced on soil health. Exhaustive efforts have been made to explore soil microbial diversity of indigenous community, their distribution and behavior in soil habitats. Use of organic manures alone or in combination with chemical fertilizers, helps in improving physico-chemical properties of the soil and also improves the efficient utilization of applied fertilizers.

**Keyword:** *Emblica officinalis*, JivaAmrit, Vermicompost, Azotobacter, Rhizogold, Bio-N-Plus.

### Introduction

Aonla (*Emblica officinalis*) is an economically important commercial fruit plants species belonging to family Euphorbiaceae. The plant is winter hardy and can be grown successfully in arid and semi-arid regions and in soils with high pH and on waste lands (Prasad and Banker, 1998). Indian Gooseberry or Aonla, *Emblica officinalis* (*Phyllanthusemblica*) is an important horticulture crop of India. It is a moderate sized deciduous tree. The fruit, bark and leaves are used in dyeing and tanning. It has aroused good deal of interest among the scientific

workers because it is one of the richest natural sources of vitamin 'C'. The fruits are major constituent of 'Chyavanprash' and 'Trifla'. The other uses of fruits are in pickle, marmalade, jam and sauce preparation. The use of bio- fertilizers in enhancing plant growth and yield has gained momentum in recent years because of higher cost and hazardous effect of chemical fertilizers. Nitrogen fixing bacteria and arbuscularmycorrhizal fungi were found to enhance growth and production of various fruit plants significantly (Khanizadehet al., 1995; DibutAlverzet al., 1996; Ghazi, 2006),

besides improving the microbiological activity in the rhizosphere (Camprubiet *al.*, 1995; Kohler *et al.*,2007).

Bio-fertilizers are the preparation containing living cells or spores, which can supply one or few plant food elements on inoculation. The farming community shows much interest in the use of bio-inoculants in a single packet as Biomix of *Azopus* or *Rhizopus*, instead of supplying it as individual organism. Several field studies on dual inoculation indicated the compatibility of N<sub>2</sub> fixing and 'P' solubilizing microorganisms. Subbarao (1982) reported the positive response of combined inoculation of N<sub>2</sub> fixing and P solubilizing microorganisms. This shows the compatibility and synergistic action among the inoculants. Similarly, the compatibility of *Rhizobium* and *phosphobacteriainoculants* was reported earlier in cowpea. Hence, a preliminary attempt had been made to study the success of *Rhizopuson* nodulation of cowpea. The chemical fertilizer no doubt increased the production but also produce many harmful effects. Therefore in developing countries like India, the use of bio-fertilizer is both economical and environmental friendly.

In order to increase the growth and yield of the crops, the farmers throughout the world are applying heavy dose of chemical fertilizers like urea, potash, super phosphate etc. It may lead to environmental pollution when the ground is water logged or the crop is not able to use the fertilizers. These fertilizers may runoff to the surface water or leaching in to the ground water. Another impact of the chemical fertilizers is it can able to lower the pH of the soil (Reynder and Viassak, 1979). Inoculation of Paddy with *Azospirillum* and *Mycorrhiza* increased the plant height and biochemical content (Kapulniket *al.*, 1981). The present work is planned to study effect of different bio-treatments on growth behavior as height, number of leaves, root length and survival of plants in nursery on *Emblicaofficinalis*. The present work was planned to study effect of

different bio-treatments on growth behavior as height, root length and survival of plants in nursery on *Emblicaofficinalis*.

## Materials and Methods

The seeds of *Emblicaofficinalis* were graded and uniform size was used for raising seedlings. Seedlings were raised in a mixture of sand: loam soil: Farm Yard Manure (1: 1: 1) in polythene bag. In order to find out suitable bio-inoculants and their combinations to achieve maximum overall growth and minimize the cost of seedling production, following treatments were given after germination. The experimental site consisted of sandy loam soil and was neutral in reaction. The water holding capacity of soil was good. The seedlings were raised in 13 × 26 cm. size polyethylene bags with a potting mixture of unsterilized sand: loam soil: farm yard manure (1:1:1). After germination in polyethylene bags 10 ml / g of broth / culture of 11 bio-treatments were inoculated in the polybags by making holes in the rhizosphere of seedlings. The treatments taken for the study were as following: T<sub>1</sub>-Control, T<sub>2</sub>-Phosphobacterium, T<sub>3</sub>-Bio P Plus, T<sub>4</sub>-Azotobacter, T<sub>5</sub>-Bio-K-Plus, T<sub>6</sub>-Bio-N-Plus, T<sub>7</sub>-JivaAmrit-liquid, T<sub>8</sub>- JivaAmrit-powder, T<sub>9</sub>-Vermicompost, T<sub>10</sub>-Azoline, T<sub>11</sub>-Rhizogold, and T<sub>12</sub>-FYM. Nursery experiment was conducted at nursery to study the effect of bio-treatments on growth behavior of Eucalyptus hybrid seedlings. The experiment was set up in a completely randomized block design with 12 treatments and three replications. All the plants were kept under identical nursery conditions and data was recorded after 90 days of application.

## Data recording

Increment in height: The plant height was measured from the base of plant to the terminal growing point of the main stem at 30, 60, 90 days after transplanting. The average plant height was expressed in centimeters.

Survival of plants: After completion of experiment, 90 days, the survival percent was also recorded for plants under all treatments including control.

Root length: After 90 days of experiment at the end, plants were randomly uprooted and root length recorded for all treatments including control. The root length measured from collar region to the tip of primary root with the help of a scale and the mean root length was expressed in centimeters.

### Results and Discussion

In general, all bio-treatments showed increased Plant height (cm), root length (cm) and survival (%) as compared to control. It is clear from Table 1 that treatments T4- Azotobacter, T6 - Bio-N-Plus, T7-Jiva Amrit-liquid, T8-Jiva Amrit-powder, and T9-Vermicompost remarkably showed very good increment in height and root length as compared to control with 11.53, 10.43, 10.97, 12.97 & 11.07 cm and 20.17, 21.27, 22.50, 20.63 & 22.27 cm respectively (Fig. 1 & 2). Similarly, the survival was also recorded after 90 days of experimentation (Table 1). It clear from the data that for most of the treatments, survival was 100 % but for control (C), it was only 80 %. For all the treatments, root length was very high as compared to control C. Soil microorganisms play an important role in soil biogeochemical processes which determine plant productivity. Successful functioning of introduced microbial bio-inoculants influenced on soil health. Exhaustive efforts have been made to explore soil microbial diversity of indigenous community, their distribution and behavior in soil habitats.

Bio-fertilizers more commonly known as microbial inoculants are artificially multiplied cultures of certain soil organisms that can improve

Soil fertility and crop productivity. The beneficial effects of *Mycorrhiza*, *Azospirillum* and *Phosphobacterium* related not only to their ability to produce antibacterial and antifungal compounds and growth regulators (Panday and

Kumar, 1989). Bio-fertilizers are environment friendly low cost agriculture input playing a significant role in improving nutrient availability to the crop plants when supplemented with chemical fertilizers. (Tilak and Singh, 1994) Field experiments carried out at different locations in India proved that the inoculation of bio-fertilizers showed better improvement in growth of the crops (SubbaRao *et al.*, 1979).

The results clearly indicate that inoculation of Rhizobium, VAM and Rhizobium + VAM significantly improve growth parameters as well as the biomass (Mishra, 2004). Best effect of these inoculations was observed in *Acacia nilotica*. Almost similar results were reported by Sharma *et al.* (1990) in case of *Acacia catechu*. Mishra (1995) has also reported good performance of *Leucaena leucocephala* after suitable inoculations on mine spoil. Better growth in case of dual inoculation has also been reported in other legumes by Kaushal (1996), Kaushik and Kaushik (1995). This increase can be attributed to better uptake of various nutrients from soil as suggested by Sharma *et al.* (1990). On the basis of present investigations and results obtained use of biofertilizers are highly recommended in agro forestry ecosystem. This is an eco-friendly and cost effective alternative to synthetic fertilizers. The effect of bioinoculants (*Frankia*, *Azospirillum* and *Phosphobacterium*) on growth performance of *Casuarinaequisetifolia* seedlings after inoculation with *Frankia* or *Azospirillum* or *Phosphobacterium* or combination of all the three inoculants after six months showed that total biomass was highest in *Frankia* + *Azospirillum* + *Phosphobacterium* (T ) inoculated seedlings (Saravanan, 2012).

The production of economic yield in any crop generally depends upon the cumulative effects of interactions among several factors such as genetic makeup of crop variety, climatic factors, mineral nutrition and cultural practices adopted. Various environmental factors such as

temperature, light, rainfall and relative humidity that prevail during different plant growth stages and development exert considerable influence on vegetative growth, seed yield and seed quality, as well as the incidence of certain pests and diseases. Application of organic manures has been a noble and traditional practice of maintaining soil health and fertility. The importance of organic manures

is, now-a-days realized because of high cost of fertilizers and their inherent capacity to supply most essential nutrients for a balanced nutrition to the crop growth. Organic nutrients generally facilitate crop rooting; improve water retention capacity and results in the even distribution of nutrients in soil profile.

**Table 1: Effect of bio-treatments on height increment and root length after 90 days**

S. No.	Treatment	Increment in height (in cm)	Root length of seedlings (in cm)	Survival (%)
1	T <sub>1</sub>	3.83	11.67	80
2	T <sub>2</sub>	6.90	13.50	90
3	T <sub>3</sub>	4.53	12.23	90
4	T <sub>4</sub>	11.53	20.17	100
5	T <sub>5</sub>	7.93	15.63	80
6	T <sub>6</sub>	10.43	21.27	90
7	T <sub>7</sub>	10.97	22.50	90
8	T <sub>8</sub>	12.97	20.63	80
9	T <sub>9</sub>	11.07	22.27	100
10	T <sub>10</sub>	10.30	15.33	100
11	T <sub>11</sub>	10.13	18.47	80
12	T <sub>12</sub>	5.03	15.43	80




Fig. 1 Effect of bio-treatments on height increment and root length after 90 days

**Treatments :**

T1►FYM T2►Phosphobacterium T3►Bio-P-Plus T4►Azotobacter T5►Bio-K-Plus T6►Bio-N-Plus T7►Jiva Amrit-liquid T8►Jiva Amrit-powder T9►Vermicompost T10►Azoline T11►Rhizogold T12►control



*Emblica officinalis* seedlings in the nursery at the initial stage of experimentation



Growth of seedlings of *Emblica officinalis* after 90 days of experimentation



T<sub>1</sub>, 2 & 3



T<sub>4</sub>, 5 & 6



T<sub>7</sub>, 8 & 9



T<sub>10</sub>, 11 & 12

Growth of seedlings of *Emblica officinalis* after 90 days of experimentation

Fig. 2

## Conclusion

India has made spectacular breakthrough in production and consumption of fertilizers during the last four decades. But consumption of chemical fertilizers will be quite a limiting factor for increasing agriculture production in future. Because of escalating energy cost, chemical fertilizers are not available at affordable prices to farmers. Moreover, the unbalanced and continuous use of chemical fertilizers is leading to reduction in crop yields and in imbalance of nutrients in the soil which has adverse effect on soil health.

Although, chemical fertilizers are playing a crucial role to meet the nutrient

requirement of the crop, persistent nutrient depletion is posing a greater threat to the sustainable agriculture. Therefore, there is an urgent need to reduce the usage of chemical fertilizers and in turn increase in the usage of organics which are needed to check the yield and quality levels. The aforesaid consequences have paved way to grow tomato using different organic manures and bio-fertilizer. Use of organic manures alone or in combination with chemical fertilizers, helps in improving physico-chemical properties of the soil and also improves the efficient utilization of applied fertilizers.

## References

- Camprubi, A., Calvet, C. and Estaun, V. (1995). Growth enhancement of Citrus reshni after inoculation with *Glomus intraradices* and *Trichoderma aureoviride* and associated effects on microbial population and enzyme activity in potting mixes. *Plant Soil*, 173: 223-38.
- Dibut Alverz, B., Rodriguez Nodals, A., Perez, A. and Martinez Viera, R. (1996). The effect of Azotoryzasduble function on banana (*Musa sp.*) experimental condition. *Infomusa*, 5: 20-23.
- Ghazi, N.A.K. (2006). Nursery inoculation of tomato with arbuscularmycorrhizal fungi and subsequent performance under irrigation with sterile water. *Sci. Hort.* 109: 1-7.
- Kaushal R., (1996). Studies on natural nodulation in various leguminous trees and influence of dual (*Rhizobium* and VAM) inoculation on growth of *Albizzialesbek*. Ph.D thesis UHF Nauni – Solan.
- Kaushik, J.C and Kaushik. N. (1995). Interaction between VAM and *Rhizobium* and their influence on *Albizzialesbek* Mycorrhizae: Biofertilizer for the Future. TERI, New Delhi.
- Khanizadeh, S., Hamel, C., Kianmehr, H., Buszard, D. and Smith, D.L. (1995). Effect of three arbuscularmycorrhizal fungus species and phosphorus on productivity and vegetative growth of three strawberry cultivars. *J. Plant Nutri.* 18: 1073-79.
- Kohler, J., Caravaca, F., Carrasco, L. and Rolden, A. (2007). Interactions between a plant growth-promoting rhizobacterium, an AM fungus and phosphate-solubilizing fungus in the rhizosphere of *Lactuca sativa*. *App. Soil Ecol.* 35: 480-87.
- Mishra, M.M., Kapoor, K.K., Chander, K., and Laura, R.D. (1991). Sustainable agriculture, The role of integrated nutrient management: A review, *Intern. J. Trop Agri.* 9(3) 155: 173.
- Mishra, P.K., (1995) Amendment of Mohulbani coal mine dump with water hyacinth and effect of VAM and *Rhizobium* on growth of *Leucaena leucocephala*. Proceeding First world mining congress, New Delhi. 875 -879.
- Pandey, A., and Kumar, S. (1989). Potential of *Azotobacter* and *Azospirillum* as biofertilizers for upland agriculture - A review *J. Sci. Ind. Res.* 48 (3): 134-144.

- Prasad, R.N. and Banker, G.J. 1998. Determination of nutrient standards of Aonla (*Emblica officinalis*) cultivars under arid conditions. *Ann. Arid Zone*, 32: 125-26.
- Reynders, S.M., and Vlassak, R.M. (1979). A survey on the occurrence of nitrogen fixing *spirillum* in Egypt. *Biotechnol. Newsletter*, 5:101-102.
- Sharma, R.P., Niranjana P and Banwari Lal (1990). Interaction between Rhizobium and mycorrhizal fungi and their stimulator effect on *Acacia nilotica* current trends in Mycorrhizal Research 259-272.
- Subba Rao, N. S. (1982). Advances in agricultural microbiology, agris.fao.org
- Saravanan, T.S., Rajendran, K. and K. Santhaguru (2012). Selection of Suitable Biofertilizers for Production of Quality Seedlings of *Casuarina Equisetifolia* (Forst.) Using Decomposed Coir Pith Compost in Root Trainers, *Asian J. Exp. Biol. Sci.* 3(4) 754.
- Subba Rao, N.S., Tilak, K.V.B.R., Singh, C.S., and Lakshmikumari, M. (1979). Response of few economic species of graminaceous plants to inoculation with *Azospirillum brasilense*. *Curr. Sci.* 48(3): 133-134.
- Tilak, K.V.B.R., and Geeta Singh (1994). Bio-fertilizers Research Gaps and future needs. *Fertilizer News*. 39 (4): 11-17.

### Acknowledgements

The authors are grateful to Council of Science and Technology, Uttar Pradesh for providing financial support to the project under which the research work was carried out. Special thanks are also due to Head, FRCER, Prayagraj (Indian Council of Forestry Research & Education, Dehradun) for providing constant guidance and encouragement throughout the work.

### CITATION OF THIS ARTICLE

Srivastava Anubha, Singh Rambir, Tomar Anita, Shukla Hari Om & Agarwal Yogesh Kumar Assessment of bio-treatments on growth performance of *Emblica officinalis* in nursery, *Int. J. Agriworld*, Vol. 1 [1] August 2020: 1-7.